

GM Science Review

Second Report

**An open review of the science relevant to GM crops and food
based on interests and concerns of the public**

Prepared by the GM Science Review Panel

(January 2004)

BIBLIOGRAPHY

The details of published papers and reports referred to in this Second Report, and which have already been included in the bibliography of our First Report, are not repeated here.

- ACNFP (2000) *Advisory Committee on Novel Foods and Processes: Annual Report 1999*. Ministry of Agriculture, Fisheries and Food/Department of Health, Crown Copyright, PB 4869, Appendix VIII, 73-76. <http://www.foodstandards.gov.uk/multimedia/pdfs/acnfp1999>
- Agrawal AA & Kotanen PM (2003) Herbivores and the success of exotic plants: a phylogenetically controlled experiment. *Ecol. Lett.* **6**, 712-715.
- Bais HP, Vepachedu R, Gilroy S, Callaway RM & Vivanco JM (2003) Alleopathy and exotic plant invasion: from molecules and genes to species interactions. *Science*, **301**, 1377-1380.
- Ballas N, Broido S, Soreq H & Loyter A (1989) Efficient functioning of plant promoters and poly(A) sites in *Xenopus* oocytes. *Nucl. Acids Res.* **17**, 7891-903.
- Ballester V, Granero G, Tabashnik BE, Malvar T & Ferre J (1999) Integrative mode for binding of *Bacillus thuringiensis* toxins in susceptible and resistant larvae for the diamondback moth (*Plutella xylostella*). *Appl. Env. Microb.* **65**, 1413-1419.
- Bauer F, Hertel C & Hammes WP (1999) Transformation of *Escherichia coli* in foodstuffs. *Syst. Appl. Microbiol.* **22**, 161-8.
- Bennetzen JL & Ma JX (2003) The genetic colinearity of rice and other cereals based on genomic sequence analysis. *Current Opinion in Plant Biology*, **6**, 128-133.
- Bergelson J, Purrington CB & Wichmann G (1998) Male promiscuity is increased in transgenic plants. *Nature*, **395**, 25.
- Bernstein JA, Bernstein IL, Bucchini L, Goldman LR, Hamilton RG, Lehrer S, Rubin C & Sampson HA (2003). Clinical and laboratory investigation of allergy to genetically modified foods. *Environmental Health Perspectives* **111**, 1114-1121.
- Betz FS, Hammond BG & Fuchs RL (2000) Safety and advantages of *Bacillus thuringiensis* protected plants to control insect pests. *Regul. Toxicol. Pharmacol.* **32**, 156-173.
- Breitbart M, Hewson I, Felts B, Mahaffy JM, Nulton J, Salamon P & Rohwer F (2003) Metagenomic analyses of an uncultured viral community from human feces. *J. Bacteriol.* **185**, 6220-6223.
- Breitzger P & Tonks D (2003) Inheritance and expression of transgenes in Barley. *Crop Science*, **43**, 4-12.
- Brookes G (2002) The farm level impact of using *Bt* maize in Spain. Brookes West, Canterbury, 23pp.
- Brooks DR, Bohan DA, Champion GT, Haughton AJ, Hawes C, Heard MS, Clark SJ, Dewar AM, Firbank LG, Perry JN, Rothery P, Scott RJ, Woiwod IP, Birchall C, Skellern MP, Walker JH, Baker P, Bell D, Browne EL, Dewar AJG, Fairfax CM, Garner BH, Haylock LA, Horne SL, Hulmes SE, Mason NS, Norton LR, Nuttall P, Randle Z, Rossall MJ, Sands RJN, Singer EJ & Walker MJ (2003) Invertebrate responses to the management of genetically modified herbicide-tolerant and conventional spring crops. I. Soil-surface-active invertebrates. *Phil. Trans, R. Soc. Lond. B* **358**, 1847-1862.
- Burke C, Yu XB, Marchitelli L, Davis EA & Ackerman S (1990) Transcription factor IIA of wheat and human function similarly with plant and animal viral promoters. *Nucl. Acids Res.* **18**, 3611-3620.
- Burke JM & Rieseberg LH (2003) The fitness effects of transgenic disease resistance in wild sunflowers. *Science*, **300**, 1250.
- Burke M (2003) GM crops: Effects on wildlife. ISBN 0-85521-035-4
- Champion GT, May MJ, Bennett S, Brooks DR, Clark SJ, Daniels RE, Firbank LG, Haughton AJ, Hawes C, Heard MS, Perry JN, Randle Z, Rossall MJ, Rothery P, Skellern MP, Scott RJ, Squire GR & Thomas MR (2003) Crop management and agronomic context of the Farm Scale

- Evaluations of genetically modified herbicide-tolerant crops. *Phil. Trans, R. Soc. Lond. B* **358**, 1801-1818.
- Chen ZL, Gu H, Li Y, Su Y, Wu P, Jiang Z, Ming X, Tian J, Pan N & Qu LJ** (2003) Safety assessment for genetically modified sweet pepper and tomato. *Toxicology*, **188**, 297-307.
- Chowdhury EH, Kuribara H, Hino A, Sultana P, Mikami O, Shimada N, Guruge KS, Saito M & Nakajima Y** (2003) Detection of corn intrinsic and recombinant DNA fragments and Cry1Ab protein in the gastrointestinal contents of pigs fed genetically modified corn Bt11. *J. Animal Sci.* **81**, 2546-2551.
- Cooper M, Dultsev FN, Minson T, Ostanin VP, Abell C & Klenerman D** (2001) Direct and sensitive detection of a human virus by rupture event scanning. *Nature Biotechnology*, **19**, 833-837.
- Defra** (2003a) Quantifying landscape-scale gene flow in oilseed rape. Research Report RG0216. <http://www.defra.gov.uk/environment/gm/research/epg-rg0216.htm>
- Defra** (2003b) Gene flow monitoring from the GM crop FSE sites: monitoring gene flow from the GM crop to non-GM equivalent crops in the vicinity. Part 1: forage maize. Research Report EPG 1/5/138. <http://www.defra.gov.uk/environment/gm/research/epg-1-5-138.htm>
- Defra** (2003c) The potential for oilseed rape feral (volunteer) weeds to cause impurities in later oilseed rape crops. Research Report RG0114. <http://www.defra.gov.uk/environment/gm/research/epg-rg0114.htm>
- Defra** (2003d) Modelling the effects on farmland food webs of herbicide and insecticide management in the agricultural ecosystem. Research Report EPG 1/5/188. <http://www.defra.gov.uk/environment/gm/research/epg-1-5-188.htm>
- Dewar AM, May MJ, Woiwood IP, Haylock LA, Champion GT, Garner BH, Sands RJN, Qi AM & Pigeon JD** (2003) A novel approach to the use of genetically tolerant crops for environmental benefit. *Proc. Roy. Soc. Lond. B.* **270**, 335-340.
- Dudek BR, Hammond BG, Nemeth MA, Lemen JK & Astwood JD** (2002) 13-week feeding study in rats fed grain from YieldGard® Event MON 810 corn. *Toxicologist* **66**(1S) Abstract # 930.
- English L & Slatin SL** (1992) Mode of action of delta-endotoxin from *Bacillus thuringiensis*: a comparison with other bacterial toxins. *Insect Biochem. Molec. Biol.* **22**, 1-7.
- EPA** (1998) EPA Registration Eligibility Decision (RED) *Bacillus thuringiensis*. EPA 738-R-98-004, March 1998.
- Ewen S W B & Pusztai A** (1999) The Lancet **354**, 1353-1354.
- Falk BW & Bruening G** (1994) Will transgenic crops generate new viruses and new diseases? *Science*, **263**, 1395-1396.
- Fares NH & El-Sayed AK** (1998) Fine structural changes in the ileum of mice fed on delta-endotoxin-treated potatoes and transgenic potatoes. *Natural. Toxins*, **6**, 219-233.
- Federici BA** (2002) Case study: Bt crops – a novel mode of insect control. In: Genetically Modified Crops. Assessing Safety. Ed: Atherton KT. Taylor and Francis New York, **22**, 164-200.
- Firbank LG & Forcella F** (2000) Genetically modified crops and farmland biodiversity. *Science*, **289**, 1481-1482.
- Fraser P & Grosveld F** (1998) Locus control regions, chromatin activation and transcription. *Curr. Opin. Cell Biol.* **10**, 361-365.
- Freckleton RP, Stephens PA, Sutherland WJ & Watkinson AR** (2003) Amelioration of biodiversity impacts of generically modified crops: predicting transient versus long-term effect. *Proc. Roy. Soc. Lond. B.* Published online 03pb0882.2.
- Fu H & Dooner HK** (2002) Intraspecific violation of genetic colinearity and its implications in maize. *Proc. Natl. Acad. Sci. USA*, **99**, 9573-9578.
- Germolec DR, Kimber I, Goldman L & Selgrade MJ** (2003) Key issues for the assessment of the allergenic potential of genetically modified foods: Breakout group reports. *Env. Health Perspectives*, **111**(8), 1131-1139.
- Goff et al.** (2002) A draft sequence of the rice genome (*Oryza sativa* L. ssp. *japonica*). *Science*, **5**, 296: 79-92.
- Gould F, Blair N, Reid M, Rennie TL, Lopes J & Micinski S** (2002) *Bacillus thuringiensis* resistance management: Stable isotope assessment of alternate host use by *Helicoverpa zea*. *Proceedings of the National Academy of Sciences*, Published online, doi:10.1073/pnas.242382499 (2002).

- Gregory PJ, Ingram JSI, Andersson R, Betts RA, Brovkin V, Chase TN, Grace PR, Gray AJ, Hamilton N, Hardy TB, Howdena SM, Jenkins A, Meybeck M, Olsson M, Ortiz-Monasterio a I, Palm CA, Payne T, Rummukainen M, Schulze RE, Thiema M, Valentin C, Wilkinson MJ.** (2002). Environmental consequences of alternative practices for intensifying crop production. *Agriculture, Ecosystems and Environment*. **88**, 279-290
- Haughton AJ, Champion GT, Hawes C, Heard MS, Brooks DR, Bohan DA, Clark SJ, Dewar AM, Firbank LG, Osborne JL, Perry JN, Rothery P, Roy DB, Scott RJ, Woiwod IP, Birchall C, Skellern MP, Walker JH, Baker P, Browne EL, Dewar AJG, Garner BH, Haylock LA, Horne SL & Mason NS** (2003) Invertebrate responses to the management of genetically modified herbicide-tolerant and conventional spring crops. II. Within-field epigeal and aerial arthropods. *Phil. Trans, R. Soc. Lond. B* **358**, 1863-1877.
- Hawes C, Haughton AJ, Osborne JL, Roy DB, Clark SJ, Perry JN, Rothery P, Bohan DA, Brooks DR, Champion GT, Dewar AM, Heard MS, Woiwod IP, Daniels RE, Young MW, Parish AM, Scott RJ, Firbank LG & Squire GR** (2003) Responses of plants and invertebrate trophic groups to contrasting herbicide regimes in the Farm Scale Evaluations of genetically modified herbicide-tolerant crops. *Phil. Trans, R. Soc. Lond. B* **358**, 1899-1913.
- Heard MS, Hawes C, Champion GT, Clark SJ, Firbank LG, Haughton AJ, Parish AM, Perry JN, Rothery P, Roy DB, Scott RJ, Skellern MP, Squire GR & Hill MO** (2003b) Weeds in fields with contrasting conventional and genetically modified herbicide-tolerant crops. II. Effects on individual species. *Phil. Trans, R. Soc. Lond. B* **358**, 1833-1846.
- Hin CJA, Schenkelaars P & Pak GA** (2001) *Agronomic and environmental impacts of the commercial cultivation of glyphosate tolerant soybean in the USA*. Utrecht: Centre for Agriculture and Environment (CLM).
- Ho MW & Cummins J** (2002) What's wrong with GMOs? *Science in Society*, **16**, 11-27.
- Ho M-W, Ryan A & Cummins J** (1999) Cauliflower mosaic viral promoter – a recipe for disaster? *Microb. Ecol. Health Dis.* **11**, 194-197.
- Hofmann C, Luthy P, Hutter R & Pliska V** (1988) Binding of the delta endotoxin from *Bacillus thuringiensis* to brush-border membrane vesicles of the cabbage butterfly (*Pieris brassicae*). *Eur. J. Biochem.* **173**, 85-91.
- IPCS** (1999) *IPCS International Programme on Chemical Safety - Environmental Health Criteria 217: Microbial pest control agent Bacillus thuringiensis*. World Health Organization, Geneva.
- Jackson AL, Bartz SR, Schelter J, Kobayashi SV, Burchard J, Mao M, Li B, Cavet G & Linsley PS** (2003) Expression profiling reveals off-target gene regulation by RNAi. *Nature Biotechnology*, **21**, 635-637, Brief Communications.
- Jank B & Haslberger AG** (2003) Improved evaluation of potential allergens in GM food *Trends in Biotech.* **21**(6), 249-250.
- Joubert GD, Venter MJ, Theron GC, Swanepoel A, Eulitz EG, Schroder HF & Macaskill P** (2001) South African experience with *Bt* cotton. *Integrated Crop Management*, Papers presented at a technical seminar at the 60th Plenary Meeting of the International Cotton Advisory Committee, September 16-21.
http://www.icac.org/icac/cotton_info/tis/biotech/documents/techsem/SAexperience_tis01.pdf
- Kashkush K, Feldman M & Levy AA** (2002) Gene loss, silencing and activation in a newly synthesized wheat allotetraploid. *Genetics*, **160**(4), 1651-1659.
- Kleter GA & Peijnenburg AACM** (2002) Screening of transgenic proteins expressed in transgenic food crops for the presence of short amino acid sequences identical to potential, IgE-binding linear epitopes of allergens. *BMC Structural Biology*, **2**, 8-19.
- Kleter GA & Peijnenburg AACM** (2003) Presence of potential allergy-related linear epitopes in novel proteins from conventional crops and the implication for the safety assessment of these crops with respect to the current testing of genetically modified crops. *Plant Biotechnology*, **1**, 371-380.
- Labra M et al.** (2001) Genomic changes in transgenic rice (*Oryza sativa* L.) plants produced by infecting calli with *Agrobacterium tumefaciens*. *Plant Cell Rep.* **20**, 325-330.
- Lin H-X, Rubio L, Smythe A, Jiminez M & Falk BW** (2003) Genetic diversity and biological variation among California isolates of Cucumber mosaic virus. *J. Gen. Virology*, **84**, 249-258.
- Lufkin T** (1997) Transcriptional regulation of vertebrate Hox genes during embryogenesis. *Crit. Rev. Eukaryot. Gene Expr.* **7**(3), 195-213.

- Lukens L, Zoub F, Lydiate D, Parkin I & Osborn T** (2003) Comparison of a Brassica oleracea genetic map with the genome of Arabidopsis thaliana. *Genetics*, **164**, 359-372.
- Mann RS** (1997) Why are Hox genes clustered? *Bio.Essays*, **19**, 661-664.
- Mayer K, Murphy G, Tarchini R, Wambutt R, Volckaert G, Pohl T, Dusterhoft A, Stiekema W, Entian K-D, Terryn N, Lemcke K, Haase D, Hall CR, van Dodeweerd A-M, Tingey SV, Mewes HW, Bevan MW & Bancroft I** (2001) Conservation of microstructure between a sequenced region of the genome of rice and multiple segments of the genome of Arabidopsis thaliana. *Genome Res.* **11**(7), 1167-1174.
- McClintock JT, Schaffer CR & Sjoblad RD** (1995) A comparative review of the mammalian toxicity of *Bacillus thuringiensis*-based pesticides. *Pestic. Sci.* **45**, 95-105.
- Mead PS, Slutsker L, Dietz V, McCaig LF, Bresee JS, Shapiro C, Griffin PM & Tauxe RV** (1999) Food-related illness and death in the United States. *Emerging Infectious Diseases*, **5**, 607-625.
- Murphy DJ** (2002) Biotechnology and the improvement of oil crops – genes, dreams and realities. *Pytochemistry Reviews*, **1**, 67-77.
- Noteborn HPJM, Rienenmann-Ploum ME, van den Berg JHJ, Alink GM, Zolla L & Kuiper HA** (1993) Food safety of transgenic tomatoes expressing the insecticidal crystal protein Cry1Ab from *Bacillus thuringiensis* and the marker enzyme APH(3')II. *Med. Fac. Landbouww. Univ. Gent.* **58/4b**.
- Noteborn HPJM, Rienenmann-Ploum ME, van den Berg JHJ, Alink GM, Zolla L Reynaerts A, Pensa M, & Kuiper HA** (1995) Safety assessment of the *Bacillus thuringiensis* insecticidal crystal protein Cry1A(b) expressed in transgenic tomatoes. In: American Chemical Society Series 605. Eds: Engel KH, Takeoka GR & Teranishi R Washington, 134-147.
- Perry JN, Rothery P, Clark SJ, Heard MS & Hawes C** (2003) Design, analysis and statistical power of the Farm-Scale Evaluations of genetically modified herbicide-tolerant crops. *J. Appl. Ecol.* **40**, 17-31.
- Pretty J** (2001) The rapid emergence of genetic modification in world agriculture: contested risks and benefits. *Environmental Conservation*, **28** (3), 248-262.
- Pusztai A, Bardocz S & Ewen SWB** (2003) GM Foods: Potential human health effects. 'Ch. 16, in *Food Safety: Contaminants and Toxicants*. Ed. D'Mello JPF, CABI publishing.
- Quemada H, L'Hostis B, Gonsalves D, Reardon IM, Henrikson R, Hiebert EL, Sieu LC & Slightom JL** (1990) The nucleotide sequences of the 3'-terminal regions of papaya ringspot strains W and P. *Journal of General Virology*, **71**, 203-210.
- Quist D & Chapela IH** (2001) Transgenic DNA introgressed into traditional maize landraces in Oaxaca, Mexico. *Nature*, **414**, 541-543.
- Read MA & Ball JG** (1999a) *The control of weeds with glufosinate ammonium in genetically modified crops of forage maize in the UK*. Proceedings of BCPC Conference, Weeds 1999, 847-852, British Crop Protection Council.
- Read MA & Ball JG** (1999b) Control of weeds in genetically modified crops of winter and spring oilseed rape with glufosinate ammonium in the UK. *Aspects of Applied Biology*, **55**, 27-33.
- Read MA & Bush MN** (1998) Control of weeds in genetically modified sugar beet with glufosinate ammonium in the UK. *Aspects of Applied Biology*, **52**, 401-406.
- Romeis J, Battini M & Bigler F** (2003) Transgenic wheat with enhanced fungal resistance causes no effects on *Folsomia candida* (Collembola: Isotomidae). *Pedobiologia*, **47**, 141-147.
- Roy DB, Bohan DA, Haughton AJ, Hill MO, Osborne JL, Clark SJ, Perry JN, Rothery P, Scott RJ, Brooks DR, Champion GT, Hawes C, Heard MS & Firbank LG** (2003) Invertebrates and vegetation of field margins adjacent to crops subject to contrasting herbicide regimes in the Farm Scale Evaluations of genetically modified herbicide-tolerant crops. *Phil. Trans. R. Soc. Lond. B* **358**, 1879-1898.
- Royal Society** (1999) *Review of data on possible toxicity of GM potatoes*. The Royal Society. Document 11/99. <http://www.royalsoc.ac.uk/gmplants/>
- Sayed AH, Cerda H & Wright DJ** (2003) Could *Bt* transgenic crops have nutritionally favourable effects on resistant insects? *Ecology Letters*, **6**, 167-169.

- Schnepf E, Crickmore N, Van Rie J, Lereclus D, Baum J, Feitelson J, Zeigler DR & Dean DH** (1998) *Bacillus thuringiensis* and its pesticidal crystal proteins. *Microbiol. Molec. Biol. Rev.* **62**, 775-806.
- Shimada N, Kim YS, Miyamoto K, Yoshioka M & Murata H** (2003) Effects of *Bacillus thuringiensis* Cry1Ab toxic on mammalian cells. *J. Vet. Med. Sci.* **65**(2), 187-191.
- Siegel JP** (2001) Minireview: The mammalian safety of *Bacillus thuringiensis*-based insecticides. *J. Invert. Pathol.* **77**, 13-21.
- Snow AA, Pilson D, Rieseberg LH, Paulsen MJ, Pleskac N, Reagon MR, Wolf DE & Selbo SM** (2003) A *Bt* transgene reduces herbivory and enhances fecundity in wild sunflowers. *Ecological Applications*, **13**, 279-286.
- Spok A, Hofer H, Valenta R, Kienzl-Plochberger K, Lehner P & Gaugitsch H** (2002) *Toxikologie und allergologie von GVO-produkten: Empfehlungen zur Standardisierung der Sicherheitsbewertung von gentechnisch veränderten Pflanzen auf Basis der Richtlinie 90/220/EWG (2001/18/EG)* Federal Environment Agency Monographien Band 109 Wien, Austria. http://www.umweltbundesamt.at/publikationsdetail.html?&pub_id=1066
- Squire GR, Brooks DR, Bohan DA, Champion GT, Daniels RE, Haughton AJ, Hawes C, Heard MS, Hill MO, May MJ, Osborne JL, Perry JN, Roy DB, Woiwod IP & Firbank LG** (2003) On the rationale and interpretation of the Farm Scale Evaluations of genetically modified herbicide-tolerant crops. *Phil. Trans, R. Soc. Lond. B* **358**, 1779-1799.
- Stewart CN, Halfhill MD & Warwick SI** (2003) Transgenic introgression from genetically modified crops to their wild relatives. *Nature Reviews Genetics*, **4**, 806-817.
- Strandberg B & Pedersen, MB** (2002) Biodiversity in glyphosate tolerant fodder beet fields - timing of herbicide application. Silkeborg: National Environmental Research Institute.
- Tepper D, Garcia-Gonzales R, Mansouri H, Seruga M, Message B, Leach F, & Perica MC** (2003) Homology-dependent DNA transfer from plants to a soil bacterium under laboratory conditions: implications in evolution and horizontal gene transfer. *Transgenic Research* **12**, 425-437.
- Thomas WE & Ellar DJ** (1983) Mechanism of action of *Bacillus thuringiensis* var. *israelensis* insecticidal δ -endotoxin. *FEBS Lett.* **154**, 362-368.
- Toivanen P, Vaahtovuori J & Eerola E** (2001) Influence of major histocompatibility complex on bacterial composition fecal flora. *Infection and Immunity*, **69**, 2372-2377.
- Van Mellaert H, Botterman J, Van Rie J & Joos H** (1999) Recombinant plant expressing non-competitively binding insecticidal crystal proteins. U.S. Patent 5,866,784.
- Van Slageren MW** (1994) Wild wheats: a monograph of *Aegilops L.* and *Amblyopyrum* (Jaub and Spach) Eig (Poaceae). Wageningen Agriculture University Papers, (7), 513pp.
- Vazquez-Padron RI, Gonzales-Cabrera J, Garcia-Tovar CB, Neri-Bazan L, Lopez-Revilla R, Hernandez M, Moreno-Fierro L & de la Riva GA** (2000) Cry1Ac protoxin from *Bacillus thuringiensis* sp. *kurstaki* HD-73 binds to surface proteins in the mouse small intestine. *Biochem. Biophys. Res. Commun.* **271**, 54-58.
- Vazquez-Padron RI, Moreno-Fierros L, Neri-Bazan L, de la Riva GA & Lopez-Revilla R** (1999) Intragastric and intraperitoneal administration of Cry1Ac protoxin from *Bacillus thuringiensis* induces systemic and mucosal antibody responses in mice. *Life Sci.* **64**, 1897-1912.
- Veraksa A, Del Campo M & McGinnis W** (2000) Developmental patterning genes and their conserved functions: From model organisms to humans. *Mol. Genet. Metab.* **69**, 85-100.
- Vlasák J, Šmahel M, Pavlík A, Pavingerová D & Bříza J** (2003) Comparison of hCMV immediate early and CaMV 35S promoters in both plant and human cells. *J. Biotechnol.* **103**, 197-202.
- Wang G et al.** (1996) Poplar (*Populus nigra L.*) plants transformed with a *Bacillus thuringiensis* toxin gene: insecticidal activity and genomic analysis. *Transgenic Res.* **5**, 289-301.
- Warwick SI, Sumard MJ, Légère A, Beckie HJ, Zhu B, Mason P, Seguin-Swartz G, & Stewart CN** (2003) Hybridization between transgenic *Brassica napus L.* and its wild relatives: *Brassica rapa L.*, *Raphanus raphanistrum L.*, *Sinapis arvensis L.*, and *Erucastrum gallicum* (Willd.). *OE Schulz. Theor. Appl. Genet.* **107**, 528-539.
- Watkinson AR, Freckleton RP, Robinson RA & Sutherland WJ** (2000) Predictions of biodiversity response to genetically modified herbicide-tolerant crops. *Science* **289**, 1554-1557.
- Wevers JDA** (1998) Agronomic and environmental aspects of herbicide-resistant sugar beet in the Netherlands. *Aspects of Applied Biology*, **52**, 393-399.

- Wilkinson MJ, Elliott LJ, Allainguillaume J, Shaw MW, Norris C, Welters R, Alexander M, Sweet J, & Mason DC** (2003) Hybridization between *Brassica napus* and *B. rapa* on a national scale in the United Kingdom. *Science*, **302**, 457-459.
- Willamson M** (1996). *Biological Invasions*. Chapman & Hall.
- Yu *et al.*** (2002) A draft sequence of the rice genome (*Oryza sativa* L. ssp. *indica*). *Science*, **5**, 296: 92-100.

GLOSSARY OF TERMS AND ABBREVIATIONS

This glossary includes the glossary of scientific and technical terms for our First Report, which was posted on the GM Science Review website shortly after its publication, and abbreviations used in both the First and Second Reports.

Abiotic	Non-biological; abiotic factors include frost, heat, drought, cold and salt.
ACAF	Advisory Committee on Animal Feedingstuffs (for the UK)
ACNFP	Advisory Committee on Novel Foods and Processes (for the UK)
ACRE	Advisory Committee on Releases to the Environment (for the UK)
AEBC	Agriculture and Environment Biotechnology Commission
<i>Agrobacterium tumefaciens</i>	A naturally occurring soil bacterium that is capable of inserting DNA (genetic information) into plants. Used in agricultural biotechnology to carry transgenes into plants.
Alien species	Species introduced intentionally or unintentionally to locations beyond the native range of the species (usually taken as post-1500). Also known as non-indigenous, non-native, exotic or introduced species. See also invasiveness.
Allele	Any one of a number of alternative forms of the same gene occupying a given position on a chromosome.
Allelopathy	Interaction between two different species involving chemicals produced by one, which suppresses the growth or reproduction of the other.
Allergenic/allergens	Substances that cause an allergic reaction.
Anaphylaxis	A severe and rapid allergic systemic reaction to contact with an allergenic trigger substance. The classic form involves prior sensitisation with later re-exposure, producing symptoms via an immunological mechanism.
Antigen	A macromolecule that is recognized by antibodies or immune cells and can trigger an immune response.

Antisense techniques	Method for down regulating the expression of particular genes by putting a ‘reverse’ version or ‘mirror image’ version of the gene, or part of the gene, into a cell.
Apomixis	Biological reproduction by seeds without fertilisation, meiosis or production of gametes. The asexual production of seeds.
ARM	Antibiotic resistance marker
Atrazine	See Triazines.
BA	British Association for the Advancement of Science
Back-crossing	Crossing an individual with one of its parents or with a genetically equivalent organism. The offspring of such a cross are referred to as the backcross generation or backcross progeny.
Bacteriophage	A virus that infects bacteria. Altered forms are used as cloning vectors.
Base pairs (bp)	Two nucleotides, one on each of the opposite complementary strands of DNA or RNA, connected via hydrogen bonds are called a base pair. In DNA, adenine and thymine, as well as guanine and cytosine, can form a base pair. The length of a nucleic acid molecule is often given in terms of the number of base pairs it contains. Modified to kbp (kilobase pairs, often used to measure transgene sizes) and Mbp (megabase pairs, used to measure total genome sizes).
BBSRC	Biotechnology and Biological Sciences Research Council
Biodiversity	A measure of the range and abundance of living organisms, this includes diversity within species and between species.
Bioinformatics	The science of managing and analysing biological data using advanced computing techniques. Especially important in analysing genomic research data. See also <i>in silico</i> .
Biolistics	Method for introducing genetic material into plant cells. DNA is coated onto micro-particles (gold or tungsten) that are fired into plant tissue.
Bolters	Plants that flower prematurely; most commonly sugar beet bolters which flower in the first year after planting rather than the second year.
bp	See base pairs

Break crop	A crop grown to benefit the soil and reduce pests and pathogens.
Bromoxynil	A nitrile herbicide that inhibits photosynthesis (in photosystem II).
<i>Bt</i>	<i>Bacillus thuringiensis</i>
Canola	Oilseed rape (used in North America).
CEC	Commission for Environmental Cooperation
Chimaera	(or chimera) An organism made up of cells containing different genetic information.
Chloroplast	Specialised plastid (general term for plant cell organelles which carry non-nuclear DNA) that contains chlorophyll. They are the site of solar energy transfer and some important reactions involved in starch and sugar synthesis. Chloroplasts have their own DNA genome and this is generally inherited through the female parent.
Clearfield	A commercial herbicide tolerant system using a group of conventionally bred crops with resistance to the broad-spectrum herbicide imidazole.
Cleistogamy	Relating to a flower that does not open and is self-pollinated in the bud.
Codon	A group of three consecutive nucleotides that represent the unit of genetic coding by specifying a particular amino acid during the synthesis of polypeptides in a cell. Each codon is recognised by a transfer RNA carrying a specific amino acid, which is incorporated into a polypeptide chain during protein synthesis.
Codon usage	As there are 64 possible triplet combinations of the 4 nucleotides (4x4x4) that are present in mRNA and DNA and only 20 amino acids, there is approximately a 3-fold excess of triplet combinations available to encode these amino acids. This means that codons can be changed without altering the amino acid sequence of the protein they encode. For some amino acids the codons most commonly used to encode them differ between eukaryotes (e.g. plants) and prokaryotes (e.g. bacteria). Therefore, the codons of a transgene derived from a prokaryote may be changed so that expression of the transgene in plant cells is more efficient.

Co-existence	In this Review, co-existence refers to the simultaneous but separate cultivation of crops by different agricultural methods (e.g. conventional non-GM, GM, organic, non-food industrial and certified seed crops).
Coleoptera	Insect order that includes beetles.
CMV	Cucumber mosaic virus
Conjugation	Unidirectional transfer of plasmid DNA from one bacterium to another, involving cell-to-cell contact. The plasmid usually encodes the majority of functions necessary for its own transfer.
Conservation headlands	Crops at the edges of fields that are not treated with agrochemicals (part of qv Countryside stewardship scheme).
Containment	Growth conditions for organisms including bacteria and plants where they are not freely released in the local environment.
COT	Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment
Countryside Stewardship scheme	A Government-run scheme that makes payments to farmers and other land managers to enhance and conserve English landscapes, their wildlife and history and to help people to enjoy them.
CP	Coat protein
Crop rotation	Crops on a specific area of land are changed year by year in a planned sequence.
Cry	Cryptochrome
cv	Cultivar
Defective interfering (DI) RNA	Small RNA molecules derived from viral RNA by extensive deletion. DI RNA depends on the original virus for replication and usually reduces the level of replication of the host (helper) virus.

Defective virus	A virus that, by itself, is unable to reproduce at all, or at the level of the wild type virus when infecting its host cell, but that can grow in the presence of another virus. This other virus provides the necessary molecular machinery that the first virus lacks.
Diploid	Having two complete sets of chromosomes, most commonly one set of paternal origin and the other of maternal origin.
DNA	Deoxyribonucleic acid. The molecule that encodes genetic information. DNA is a double-stranded molecule held together by weak bonds between base pairs of nucleotides. The four nucleotides in DNA contain the bases adenine (A), guanine (G), cytosine (C), and thymine (T). In nature, base pairs form only between A and T and between G and C; thus the base sequence of each single strand can be deduced from that of its partner. The structure of DNA (double helix) was published in 1953 by Crick and Watson.
Ecosystem	The complex of a living community and its physical environment, functioning as an ecological unit in nature.
EEA	European Environment Agency
EFSA	European Food Safety Authority
Embryo rescue	A sequence of tissue culture techniques used to enable a fertilised immature embryo, resulting from an interspecific cross (between plants from different species), to continue growth and development (when it may not otherwise), until it can be regenerated into an adult plant.
Encapsulation	Packaging of viral genetic material (RNA or DNA) in a regular protective, geometric array (shell) or coat (capsid) protein molecules.
ENTRANSFOOD	European Network for Safety Assessment of Genetically Modified Food Crops
EPA	Environmental Protection Agency (for the USA)
Epidemiological monitoring	See Post-marketing monitoring
Epidemiological surveillance	See Post-marketing surveillance
Epitopes	The individual surface feature of an antigen that elicits the production of a specific antibody (monoclonal) in the course of the immune response. Each antigenic determinant, typically a

few amino acids in size, causes the synthesis of a different antibody and thus exposure to a single antigen may result in the expression of a number of antibodies (polyclonal).

ESTO	Earth Science Technology Office
EU	European Union
Eukaryote/ eukaryotic	Cell or organism with membrane-bound, structurally discrete nucleus and other well-developed subcellular compartments. Eukaryotes include all organisms except viruses, bacteria, and bluegreen bacteria (algae).
Exon	Protein-coding DNA sequence of a gene. Many eukaryotic genes are composed of a mosaic of exons and introns.
F1 hybrids	The initial hybrid generation resulting from a cross between two genetically unlike parents.
FAO	Food and Agriculture Organisation (of the United Nations)
Farm-saved seed	Retaining some seed from a harvest to sow in following years on the same farm, rather than buying in new seed produced commercially.
FDA	Food and Drug Administration (for the USA)
Fertilisation (self and cross)	Union of two gametes from opposite sexes to form a zygote. There are different categories: 1. Self-fertilisation (selfing): fusion of male and female gametes from the same individual. 2. Cross-fertilisation (crossing): fusion of male and female gametes from different individuals.
FISH	Fluorescence <i>in-situ</i> hybridisation. A method for locating particular sections of DNA on chromosomes.
Fitness	The survival value and the reproductive capability of an individual, compared to that of competitor individuals of the same or other species within a population or an environment.
Fixation	Situation in which only one allele for a given gene/locus is present in a population. This can occur as a result of direct selection where the allele delivers a greater level of fitness; because of indirect selection, where the locus is linked to a gene that is subject to direct selection; or because of genetic drift. Also: Nitrogen fixation: the process of conversion of gaseous nitrogen in the atmosphere to compounds which can be metabolized by plants, a process which can be carried out by some bacterial species.

FOE	Friends of the Earth
FSA	Foods Standards Agency (for the UK)
FSEs	GM Crop Farm-Scale Evaluations
Fusion protein	A polypeptide synthesised from a chimeric gene. The different fragments of DNA are joined so that their protein coding sequences are in the same reading frame. The resulting construct is transcribed and translated as a single gene, producing a single protein.
Gamete	Mature male or female reproductive cell.
Gene	The unit of heredity transmitted from generation to generation during sexual or asexual reproduction. The simplest gene consists of a segment of nucleic acid that encodes an individual protein or a length of RNA.
Gene construct	The DNA unit, usually including transgenes, promoters and selectable markers, which is used to make a GM plant.
Gene flow	The transfer of genes between different individuals e.g. pollen-mediated gene transfer between sexually compatible plants. Also refers to the transfer of genes from one plant population to another through seed dispersal or the movement of regenerative plant parts, (e.g. tubers), or whole plants. This Review also considers the possibility of plant genes being transferred and stably integrated into the genomes of soil and gut microbes and into viruses that infect plants (see horizontal gene flow).
Gene product	RNA and proteins.
Gene stacking	Accumulation of genes conferring different traits in one plant resulting from cross-fertilisation or transformation with several gene constructs. Also, see transgene stacking.
Genotype	The genetic constitution of an organism, as distinguished from its physical characteristics (its phenotype).
GLP	Good Laboratory Practice
Glufosinate ammonium	Used to provide post-emergence, broad spectrum control of annual grasses and broad-leaved weeds. Glufosinate ammonium can be sprayed after emergence if the crop is tolerant to it. This herbicide acts by inhibiting an enzyme that is responsible for ammonia detoxification ultimately leading to the cessation of photosynthesis. The trade names of herbicides

containing glufosinate ammonium include: Basta, Liberty, Ignite, and HOE 39866.

Glyphosate	Systemic herbicide that is used for post-emergence, broad spectrum control of annual and perennial broad-leaved and grass weeds. Can be sprayed after emergence if the crop is glyphosate tolerant. Acts by inhibiting an amino acid metabolism pathway that exists in higher plants and microorganisms, but not in animals. Inactivated on contact with clay particles in soil, and requires no hazard warning symbols on packaging. The trade names of some herbicides in which glyphosate is the active ingredient are: Roundup, Rodeo, Touchdown, and MON-0573.
GM	Genetically modified/ Genetic modification. Altering the genetic material of an organism in a way that does not occur naturally by mating and / or natural recombination.
GM derived	Products that are derived from genetically modified organisms, including products (e.g. some vegetable oils or enzymes used for making cheese) in which it is not possible to detect any DNA or protein.
GMHT	Genetically Modified Herbicide Tolerance
GMIR	Genetically Modified Insect Resistant
GMO	Genetically modified organism. An organism in which the genetic material has been altered in a way that does not occur naturally by mating and / or natural recombination.
GFP	Green fluorescence protein
GI	Gastro-intestinal
GURT	Genetic use restriction technologies. Mechanisms that prevent or restrict gene flow from GM crops, e.g. GM male sterility.
Haploid	Possessing only one copy of each chromosome. Within higher organisms, only the reproductive cells are haploid, whereas the somatic (body) cells are diploid (two copies of each chromosome) or polyploid (three or more copies of each chromosome - often found in plants).
Homologous recombination	See Recombination.
Homology	Similarity between sequences of DNA or amino acids (that make up proteins) in individuals of the same, or different species.

Horizontal gene flow	See Horizontal gene transfer.
Horizontal gene transfer (HGT)	Non-sexual, non-parental-to-offspring processes by which genetic material can sometimes transfer between organisms with distant genetic relationships.
Hybrids/hybridisation	Offspring of two genetically unlike parents. Crossing of two sexually compatible but genetically different plants.
ICSU	International Council of Scientific Unions
IFPRI	International Food Policy Research Institute
IgE	Immunoglobulin E
ILGRA	Interdepartmental Liaison Group on Risk Assessment (for the UK)
ILSI	International Life Sciences Institute
<i>In silico</i>	In a computer. In the present context, the use of data bases of DNA and protein sequence to help answer biological questions. This is a growing area of biology, as the amount of genomic and proteomic data continue to grow. See bio-informatics.
Interspecific plant competition	Competition between one plant species and another.
Introgression	Introduction of new allele(s) or gene(s) into a population from an exotic source, usually another species. This is achieved by repeated backcrossing of the initial hybrid in order to eliminate all genetic changes except for the desired new gene(s).
Intron	DNA sequence that interrupts the protein-coding sequence of a gene. An intron is transcribed into RNA but is cut out of the message before it is translated into protein.
Invasiveness (or invasive species)	Ability of an organism, particularly an alien species (qv), to spread beyond its presently established site, and become established in new locations.
<i>in vitro</i>	Outside the organism, or in an artificial environment. Applied, for example, to cells, tissues or organs cultured in glass or plastic containers.
<i>in vivo</i>	The natural conditions in which organisms reside. Refers to biological processes that take place within a living organism or cell under normal conditions.

JRC	Joint Regulatory Commission (for the European Union)
Late successional closed vegetation	The endpoint of natural change following disturbance (e.g. forest or heath).
Lepidoptera	Order of insects that includes butterflies and moths.
Linkage drag	When the selection pressure operating for, or against, a trait encoded by a gene(s) is affected by the co-inheritance of linked genes.
Macrophage	Large white blood cells that ingest foreign substances and display on their surfaces antigens that are recognised by other cells of the immune system.
Marker genes	A gene of known function, sequence and/or location, used for marker-assisted selection or for genetic studies. Marker genes conferring traits that are novel to plants are used to identify plants into which transgenic DNA has been successfully introduced e.g. genes encoding resistance to certain antibiotics.
Marker rescue	Restoration of gene function by replacement of a defective gene with a normal one through recombination.
Meiosis	Process of two consecutive cell divisions in the formation of sex cells (gametes).
Metabolomics	The large-scale study of the full complement of secondary metabolites produced by a given species in all its cells, tissues and/ or growth stages.
Microarrays	A large set of DNA molecules immobilized as a compact and orderly pattern of sub-microlitre spots onto a solid matrix (e.g. a glass slide). Used to analyse patterns of gene expression, presence of markers, or nucleotide sequences. The major advantage of microarrays is the extent to which the process of genotyping can be automated, thereby enabling large numbers of individuals to be genetically typed at many loci simultaneously. A similar approach may be used with other immobilised components (e.g. proteins) for other purposes.
Micronutrients	Components of nutrition required in relatively small quantities by organisms, e.g. vitamins or minerals.
Millieumetlat system	Scoring system is used to evaluate toxicity, mobility and persistence of pesticides.
mRNA	Messenger RNA. The RNA molecule resulting from transcription of a protein-encoding gene, following any

splicing (removal of introns). The information encoded in the mRNA molecule is translated into a polypeptide (protein) by the ribosomes.

mRNA fingerprinting

Pattern of mRNAs present in an organism under specific conditions.

Mutation breeding

Induction of heritable change(s) in the genetic constitution of a cell through alterations to its DNA, using mutagenic chemicals or radiation. Used in breeding programmes to introduce genetic differences from which new crop phenotypes with desirable traits can be selected. Can cause gross and unpredictable changes to whole chromosomes as well as to specific genetic loci. These smallest changes can involve the substitution, deletion or insertion of a single nucleotide.

Naked DNA

DNA from any source (whether or not created by recombinant DNA techniques) that has been purified and separated from the proteins that normally surround the DNA in a living organism.

NGOs

Non-Governmental Organisations

Nuclear DNA

DNA organised into chromosomes and which contains most of the genes (typically 25 000 – 50 000) that are largely responsible for the differentiation and activity of the cell. Plastids and mitochondria contain non-nuclear DNA.

Nucleotides

A subunit of DNA or RNA consisting of a nitrogenous base (adenine, guanine, thymine, or cytosine in DNA; adenine, guanine, uracil or cytosine in RNA), a phosphate molecule and a sugar molecule (deoxyribose in DNA and ribose in RNA). Thousands of nucleotides are linked through repeating sugar-phosphate bonds to form a DNA or RNA molecule.

OECD

Organisation for Economic Cooperation and Development

Open reading frames

Abbreviation: ORF. A sequence of nucleotides in a DNA or RNA molecule that has the potential to encode a peptide or protein: comprises a start triplet (ATG), followed by a series of triplets (each of which encodes an amino acid), and ending with a stop codon (TAA, TAG or TGA). The number of ORFs provides an estimate of the number of genes that could be transcribed from the DNA sequence.

ORFs

Open reading frames

Origin of replication

See Replication origin.

Pathogen

Disease-causing organism (generally microbial: bacterial, fungal or viral; but can extend to other organisms, e.g.

	nematodes).
Pest	An organism that reduces the productivity of a crop e.g. certain insects, birds and nematodes.
PCR	See Polymerase chain reaction.
Phytochemicals	Molecules characteristically found in plants.
Phytoremediation	Biological remediation (restoration) of the environment using plants.
Plasmid	Autonomously replicating extra-chromosomal circular DNA molecule, distinct from the normal bacterial genome and non-essential for cell survival under non-selective conditions. Some plasmids are capable of integrating into the host genome. A number of artificially constructed plasmids are used as gene cloning vectors (vehicles).
Plastid	Plant-specific organelles, such as chloroplasts, which carry their own DNA.
Pleiotropy	The simultaneous effect of a gene on more than one apparently unrelated trait.
PLRV	Potato leafroll virus
Pollination	Part of the process of fertilisation, in which pollen is transferred from an anther (male part) to the stigma (female part) of the same (self-pollination) or a different (cross-pollination) sexually compatible plant.
Polymerase chain reaction	Abbreviation: PCR. A method for dramatically increasing the number of copies of a specific fragment of DNA <i>in vitro</i> .
Polyploid	Three or more copies of each chromosome in a cell.
Post-marketing monitoring	The hypothesis-driven, routine collection of information after a product is on the market (i.e. widely available). For example, epidemiological monitoring involves looking for a disease condition, characteristic or state in a population.
Post-marketing surveillance	Surveillance takes a general look at trends. e.g. epidemiological surveillance is the systematic collection, collation, analysis and interpretation of health-related events occurring in populations.
Prokaryotes	Unicellular organism lacking a membrane-bound, structurally discrete nucleus and other subcellular compartments. Bacteria are examples of prokaryotes.

Promoter	A DNA sequence at the start of a gene to which RNA polymerase (an enzyme) will bind and initiates transcription/expression of a gene into messenger (or other) RNA. Genomic and subgenomic promoters also exist in RNA viruses where they initiate copying of RNA into RNA.
PROSAMO	Planned release of selected and modified organisms
Protists	A single-celled eukaryote.
Recombinant DNA technology	Set of techniques for manipulating DNA, including: the identification, modification and cloning of genes; the study of the expression of cloned genes; and the production of large quantities of gene products.
Recombinase	Class of enzymes able to alter the arrangement of DNA sequences in a site-specific way.
Recombination	Process by which progeny derive a combination of genes different from that of either parent. In higher organisms such as plants, this can occur by crossing over (breaking during meiosis of one maternal and one paternal chromosome, the exchange of corresponding sections of DNA, and the rejoining of the chromosomes). In lower organisms such as bacteria and viruses, it describes the cutting and rejoining or RNA/DNA template switching, which results in the exchange of fragments of genetic material or information between different organisms. It also describes the transfer of transgenic material between lower organisms and GM plants containing homologous sequences. Homologous recombination mediates the transfer or exchange of genetic information between homologous sections of DNA.
Refugia	Area of crops or adjacent land where there is no control of weeds or, more usually, insects, and thus provide a safe haven for them. Refugia can be placed adjacent to conventional crops sprayed with herbicides or insecticides or near GM herbicide-tolerant or insect-resistant crops to reduce the selection pressure on the insects or weeds to evolve resistance and increase local biodiversity. May be accomplished by qv: Countryside Stewardship Schemes, Conservation Headlands and Wildlife strips.
Replication origin	The nucleotide position on a DNA sequence from which DNA synthesis (replication) is initiated.
Reverse transcriptase PCR	Abbreviation: RT-PCR. RNA (mRNA) molecule(s) reverse transcribed into a DNA copy are then amplified using PCR. RT-PCR can be used to identify which genes are being

expressed in a cell (i.e. the mRNA population), whereas PCR identifies all of the genes (DNA) that are present.

RNAi	RNA interference.
RSPB	Royal Society for the Protection of Birds
Satellite RNA	Plant viruses often contain parasites of their own, referred to as satellites. Satellite RNAs are dependent on their associated (helper) virus for both replication and encapsidation.
SCIMAC	Supply chain initiative on modified agricultural crops
Secondary metabolite profiling	See Metabolomics.
Silencing	Mechanisms in a genome that repress the expression of genes; can be achieved using transgenes. Recent work involves qv RNAi.
siRNA	Small interfering RNA
SSCP	Single-strand (RNA) conformation polymorphism
SSSI	Site of Special Scientific Interest
Substantial equivalence	Used to structure the comparison of a novel food with its conventional counterpart to identify any compositional differences that then become part of a more focussed safety assessment.
Sulfonyl urea	Herbicides that block the synthesis of essential branched chain amino acids (leucine, isoleucine and valine) by inhibiting the enzyme acetolactate synthase (ALS). For this reason, these chemicals are sometimes referred to as SU/ALS herbicides or ALS herbicides.
Synergism	Phenomenon in which one virus may facilitate replication and/or increase the symptom severity of another co-infecting virus, resulting in more severe disease.
Terminator	DNA sequence just downstream of the coding segment of a gene, which is recognized by RNA polymerase as a signal to stop synthesizing mRNA.
Terminator technology	A type of GURT. Transgenic method that genetically sterilises the progeny of a planted seed.
Tillage	Ploughing or harrowing. Zero-tillage or low-till agricultural practices may be implemented.

Trait	One of the many characteristics that define an organism. The phenotype is a description of one or more traits.
Trans	Spatially separated, e.g. on opposite chromosomes.
Transencapsulation	Complete or partial encapsidation (packaging), of the genome of one virus with the coat protein of another virus.
Transformation	Uptake and integration of DNA in a cell.
Transgene stacking	Accumulation of transgenes conferring different traits in one plant. This can arise intentionally or unintentionally through cross-fertilisation or by the introduction of different traits into a GM plant variety through one or a number of successive transformation events.
Transgenic DNA/transgene	Isolated sequence of DNA stably inserted into the genome of a recipient organism.
Translation initiation signal/factor	The RNA codon (AUG) that specifies the first amino acid of a polypeptide chain. An assemblage of proteins necessary for the initiation of polypeptide synthesis from mRNA.
Transposon	DNA element that can move from one location in the genome to another, or through an RNA intermediate.
Triazines	Herbicides that inhibit photosynthesis (in particular photosystem II) e.g. atrazine.
UK	United Kingdom
UNCED	United Nations Conference on Environment and Development
Unencapsidated	Viral DNA or RNA not enclosed by a coat protein shell or capsid.
USDA	US Department of Agriculture
Vector	Small DNA molecule (plasmid, virus, bacteriophage, artificial or cut DNA molecule) that can be used to deliver DNA into a cell. Vectors must be capable of being replicated and contain cloning sites for the introduction of foreign DNA. Vector can also refer to an organism, usually an insect, which carries and transmits pathogens. Also refers to an organism, usually an insect that carries and transmits pathogens/ disease.
Vernalisation	Chilling juvenile plants for a minimum period in order to induce flowering. Some plants (e.g. sugar beet) require vernalisation to flower, but others have no such requirement.

Viroids/virusoids	Unique plant pathogenic agents, composed of infectious single-stranded low molecular weight RNAs, and no coat protein.
Volunteer	Crop plant self-propagated from a previous year's crop (e.g. from seed or tubers).
Wild type	The most frequent allele or genotype found in nature.
Wildlife strips	Edges of fields that are not planted or treated with agrochemicals (part of qv Countryside stewardship scheme).
WHO	World Health Organisation
Zygote	The result of fertilisation between two gametes. It undergoes a cycle of multiple divisions to become an embryo.

Responses to the First Report of the GM Science Review

The following organisations and individuals responded to our invitation to comment on the First Report. The great majority of responses were e-mailed¹ to us and these can be viewed on the GM Science Review website².

T Addiscott	C Creighton*
Agriculture and Biotechnology Council	E Cross
M H Arnold	J Davison
M Berwyn-Jones	A Dean
J Bond	P Dupont
Bristol Area SERA Group	J Dutton
Bristol Genetix Group ³ *	R Eburne
British Crop Protection Council	Econexus, the Five Year Freeze, Friends of the Earth, Genewatch UK, Greenpeace, the Soil Association and Dr M Antoniou
British Society of Plant Breeders Limited	
British Statutory Nature Conservation Agencies: Countryside Council for Wales, English Nature, Joint Nature Conservation Committee and Scottish Natural Heritage	P Elliot G Finch
British Trust for Ornithology	M Fuller
T Brown	H Gatt
D Burke	F A Gibson*
M Cantley	D Gledhill
A M Claparols	Green Party
J Clarke	N Grice
S Cooke	S C Harris*

¹ An asterisk against a name denotes a written response.

² <http://www.gmsciencedebate.org.uk/report/comment/default.htm>

³ Similar letters from several individuals.

G Humphreys

Independent Science Panel (ISP)

Institute of Biology on behalf of the
Biosciences Federation

Institute of Science in Society (ISIS)

S James

John Innes Centre

A Kerry-Bedell

I Langford*

U Loening

N Maclean

R Marsh

N Mullan

Munlochy Vigil

S Neil

T Nichells

E Overton

R Patterson

Penwith Organic Gardeners and Growers*

Prospect

D Reid

O Richardson*

The Scientific Alliance

Scientists for Global Responsibility

R Scullion

Swindon Friends of the Earth

Syngenta

J Tipper*

I Wade*

G White

M White*